#### Catalog # BEB-H52H3



#### Synonym

BG, MPS7

#### Source

Human beta-Glucuronidase Protein, His Tag(BEB-H52H3) is expressed from human 293 cells (HEK293). It contains AA Leu 23 - Thr 651 (Accession # <u>P08236</u>).

Predicted N-terminus: Leu 23

### **Molecular Characterization**

beta-Glucuronidase/GUSB(Leu 23 - Thr 651) P08236 Poly-his

This protein carries a polyhistidine tag at the C-terminus

The protein has a calculated MW of 74.5 kDa. The protein migrates as 75-85 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

# Endotoxin

Less than 1.0 EU per  $\mu g$  by the LAL method.

# Purity

>95% as determined by SDS-PAGE.

#### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in 50 mM Tris,150 mM NaCI,pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

# **SDS-PAGE**



Human beta-Glucuronidase Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

# **Bioactivity**

Measured by its ability to hydrolyze 4-methylumbelliferyl-beta -D-

glucuronide. The specific activity is >3500 pmol/min/µg(QC tested).

Background

>> www.acrobiosystems.com

5/12/2023



### Catalog # BEB-H52H3

Human beta -Glucuronidase (EC 3.2.1.31) encoded by the GUSB gene is a lysosomal hydrolase involved in the stepwise degradation of glucuronic acid-containing glycosaminoglycans that include heparan sulfate, chondroitin sulfate and hyaluronan. The enzyme is only active on the glucuronic acid of the non-reducing end. The native protein has been reported as a tetrameric glycoprotein composed of identical subunits.

### **Clinical and Translational Updates**

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.



5/12/2023