

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

Human c-MET (Luc) HEK293 Reporter Cell


Catalog No.	Size
CHEK-ATF144	2 × (1 vial contains ~5×10 ⁶ cells)

• Description

The Human c-MET (Luc) HEK293 Reporter Cell was engineered to express signaling response element driving luciferase expressing systems and human c-MET (Gene ID: 4233). When stimulated with human HGF protein, the HGF/c-MET interaction drives RE-mediated luminescence. Neutralization of biological effect of human HGF protein by corresponding antibody results in a decrease in luminescence.

• Application

- Screen for neutralizing antibodies blocking the stimulation of human HGF protein.



Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

• Cell Line Profile

Cell line	Human c-MET(Luc) HEK293 Reporter Cell
Host Cell	HEK293
Property	Adherent
Complete Growth Medium	DMEM + 10% FBS
Selection Marker	Puromycin (2 µg/mL)
Incubation	37°C with 5% CO ₂
Doubling Time	22-24 hours
Transduction Technique	Lentivirus

• Materials Required for Cell Culture

- DMEM medium (Gibco, Cat.No.11965-092)
- Fetal bovine serum (CellMax, Cat.No.SA211.02)
- Puromycin (InvivoGen, Cat.No.ant-pr-5b)
- Complete Growth Medium: DMEM + 10% FBS
- Culture Medium: DMEM + 10% FBS, Puromycin (2 µg/mL)
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, 430641)
- Cryogenic storage vials (SARSTEDT, 72.379.007)
- Thermostat water bath
- Centrifuge
- Luna cell counter (Logos Biosystems, LUNA- II)
- CO₂ Incubator (Thermo, 3111)
- Biological Safety Cabinet (Thermo, 1389)

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

• *Recovery*

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by spraying with 70% ethanol. All the operations from this point on should be carried out under strict aseptic conditions.
3. Transfer the vial contents to a centrifuge tube containing 4.0 mL complete growth medium and spin at approximately 1000 rpm for 5 minutes.
4. Resuspend cell pellet with 5 mL complete growth medium and transfer the cell suspension into T-75 flask containing 10-15 mL of pre-warmed complete growth medium.
5. Incubate at 37°C with 5% CO₂ incubator until the cells are ready to be split.

• *Subculture*

1. Remove and discard culture medium.
2. Wash the cells once with sterile PBS.
3. Add 2 mL of 0.25% trypsin to cell culture flask. Place the flask at 37°C for 2-3 minutes, until 90% of the cells have detached.
4. Add 6.0 to 8.0 mL of culture medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessel.
6. Incubate at 37°C with 5% CO₂ incubator.

Subcultivation Ratio: A subcultivation ratio of 1:6 to 1:10 is recommended.

Medium Renewal: Every 2 to 3 days.

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

• *Cryopreservation*

1. Remove and discard spent medium.
2. Detach cells from the cell culture flasks with 0.25% trypsin.
3. Centrifuge at 1000 rpm for 5 min at RT to pellet cells.
4. Resuspend the cell pellets with complete growth medium and count viable cells.
5. Centrifuge at 1000 rpm for 5 min at RT and resuspend cells in freezing medium to a concentration of 5×10^6 to 1×10^7 cells/mL.
6. Aliquot into cryogenic storage vials. Place vials in a programmable cooler or an insulated box placed in a -80°C freezer overnight, then transferring to liquid nitrogen storage.

• *Storage*

- **Product format:** Frozen
- **Storage conditions:** Liquid nitrogen immediately upon receipt

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

• Receptor Assay




Fig1. Expression analysis of human c-MET on Human c-MET (Luc) HEK293 Reporter Cell by FACS. Cell surface staining was performed on Human c-MET (Luc) HEK293 Reporter Cell or negative control cell using PE-labeled anti-human c-MET antibody.

• Signaling Bioassay




Fig2. Response to human HGF protein (RLU). This reporter cell was incubated with serial dilutions of human HGF protein. The EC₅₀ was approximately 0.1506 µg/mL.

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet




Fig3. Response to human HGF protein (FOLD). This reporter cell was incubated with serial dilutions of human HGF protein. The max induction fold was approximately 41.75.

• *Application*

Anti-human c-MET Neutralizing Antibody Screening




Fig4. Inhibition of human HGF protein-induced reporter activity by anti-human c-MET neutralizing antibody. This reporter cell was incubated with serial dilutions of antibodies in the presence of human HGF protein with a final concentration of 0.3 µg/mL. The EC50 of anti-human c-MET neutralizing antibody (Amivantamab) is approximately 0.07699 µg/mL.

Human c-MET (Luc) HEK293 Reporter Cell Data Sheet

- ***License Disclosure***

This reporter cell is provided for research use only. This license does not permit you to share, distribute, sell, sublicense, or otherwise make this reporter cell available for use to other laboratories, departments, research institutions, hospitals, universities, or biotech companies. The license does not permit modification of this reporter cell in any way. Inappropriate use or distribution of this reporter cell will result in revocation of the license. Modifications of this cell line, transfer to another facility, or commercial use of the cells may require a separate license and additional fees. AcroBiosystems does not warrant the suitability of this reporter cell for any particular use, and does not accept any liability in connection with the handling or use of this reporter cell.

- ***Related Products***

Products

Human EGF R (Luc) HEK293 Reporter Cell

Cat.No.

CHEK-ATF049