

# Human BMP (Luc) HEK293 Reporter Cell Data Sheet

## Human BMP (Luc) HEK293 Reporter Cell


Catalog No.	Size
CHEK-ATF188	2 × (1 vial contains ~5×10 <sup>6</sup> cells)

### • Description

The Human BMP (Luc) HEK293 Reporter Cell was engineered to express Smad signaling response element driving luciferase expressing systems. When stimulated with BMP protein, receptor-mediated signaling can drive Smad-mediated luminescence.

### • Application

- Bioactivity detection of human BMP protein



# Human BMP (Luc) HEK293 Reporter Cell Data Sheet

## • Cell Line Profile

<b>Cell line</b>	Human BMP (Luc) HEK293 Reporter Cell
<b>Host Cell</b>	HEK293
<b>Property</b>	Adherent
<b>Complete Growth Medium</b>	DMEM + 10% FBS
<b>Selection Marker</b>	Puromycin (2 µg/mL)
<b>Incubation</b>	37°C with 5% CO <sub>2</sub>
<b>Doubling Time</b>	22-24 hours
<b>Transduction Technique</b>	Lentivirus

## • Materials Required for Cell Culture

- DMEM medium (Gibco, Cat.No.11965-092)
- Fetal bovine serum (CellMax, Cat.No.SA211.02)
- Puromycin (InvivoGen, Cat.No.ant-pr-5b)
- 0.25% Trypsin-EDTA (1X), Phenol Red (Gibco, Cat.No.25200-056)
- Penicillin-Streptomycin (Gibco, Cat.No.15140-122)
- Phosphate Buffered Saline (1X) (HyClone, Cat.No.SH30256.01)
- Complete Growth Medium: DMEM + 10% FBS, 1%P/S
- Culture Medium: DMEM + 10% FBS, Puromycin (2 µg/mL), 1%P/S
- Freeze Medium: 90% FBS, 10% (V/V) DMSO
- T-75 Culture flask (Corning, 430641)
- Cryogenic storage vials (SARSTEDT, 72.379.007)
- Thermostat water bath
- Centrifuge
- Luna cell counter (Logos Biosystems, LUNA- II )
- CO<sub>2</sub> Incubator (Thermo, 3111)
- Biological Safety Cabinet (Thermo, 1389)

# Human BMP (Luc) HEK293 Reporter Cell Data Sheet

## • *Recovery*

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by spraying with 70% ethanol. All the operations from this point on should be carried out under strict aseptic conditions.
3. Transfer the vial contents to a centrifuge tube containing 4.0 mL complete growth medium and spin at approximately 1000 rpm for 5 minutes.
4. Resuspend cell pellet with 5 mL complete growth medium and transfer the cell suspension into T-75 flask containing 10-15 mL of pre-warmed complete growth medium.
5. Incubate at 37°C with 5% CO<sub>2</sub> incubator until the cells are ready to be split.

## • *Subculture*

1. Remove and discard culture medium.
2. Wash the cells once with sterile PBS.
3. Add 2 mL of 0.25% trypsin to cell culture flask. Place the flask at 37°C for 2-3 minutes, until 90% of the cells have detached.
4. Add 6.0 to 8.0 mL of culture medium and aspirate cells by gently pipetting.
5. Add appropriate aliquots of the cell suspension to new culture vessel.
6. Incubate at 37°C with 5% CO<sub>2</sub> incubator.

**Subcultivation Ratio:** A subcultivation ratio of 1:6 to 1:10 is recommended.

**Medium Renewal:** Every 2 to 3 days.

**Note:** After recovery for 1-2 generations with the complete growth medium not containing the selection marker, if the cell state is well, changing to the culture medium containing the selection marker.

## Human BMP (Luc) HEK293 Reporter Cell Data Sheet

### • *Cryopreservation*


1. Remove and discard spent medium.
2. Detach cells from the cell culture flasks with 0.25% trypsin.
3. Centrifuge at 1000 rpm for 5 min at RT to pellet cells.
4. Resuspend the cell pellets with complete growth medium and count viable cells.
5. Centrifuge at 1000 rpm for 5 min at RT and resuspend cells in freezing medium to a concentration of  $5 \times 10^6$  to  $1 \times 10^7$  cells/mL.
6. Aliquot into cryogenic storage vials. Place vials in a programmable cooler or an insulated box placed in a  $-80^\circ\text{C}$  freezer overnight, then transferring to liquid nitrogen storage.

### • *Storage*


- **Product format:** Frozen
- **Storage conditions:** Liquid nitrogen immediately upon receipt

# Human BMP (Luc) HEK293 Reporter Cell Data Sheet

• *Signaling Bioassay*




**Fig1. Response to human BMP-2 protein (RLU).** The Human BMP (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human BMP-2 protein (Cat.No.BM2-H4117). The EC50 was approximately 0.1212 µg/mL.




**Fig2. Response to human BMP-2 protein (FOLD).** The Human BMP (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human BMP-2 protein (Cat.No.BM2-H4117). The max induction fold was approximately 19.39.

# Human BMP (Luc) HEK293 Reporter Cell Data Sheet



**Fig3. Response to human BMP-4 protein (RLU).** The Human BMP (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human BMP-4 protein. The EC50 was approximately 0.2787 µg/mL.



**Fig4. Response to human BMP-4 protein (FOLD).** The Human BMP (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of human BMP-4 protein. The max induction fold was approximately 17.07.

# Human BMP (Luc) HEK293 Reporter Cell Data Sheet

## • *License Disclosure*

This reporter cell is provided for research use only. This license does not permit you to share, distribute, sell, sublicense, or otherwise make this reporter cell available for use to other laboratories, departments, research institutions, hospitals, universities, or biotech companies. The license does not permit modification of this reporter cell in any way. Inappropriate use or distribution of this reporter cell will result in revocation of the license. Modifications of this cell line, transfer to another facility, or commercial use of the cells may require a separate license and additional fees. AcroBiosystems does not warrant the suitability of this reporter cell for any particular use, and does not accept any liability in connection with the handling or use of this reporter cell.

## • *Related Products*

### Products

Human BMP-2 Protein, premium grade

### Cat.No.

BM2-H4117