

**Synonym**

HA, Hemagglutinin

**Source**

Influenza B virus [Austria/1359417/2021 (B/Victoria lineage)] HA Protein, His Tag (HAE-V52H8) is expressed from Baculovirus-Insect cells.

Predicted N-terminus: Asp 16

**Molecular Characterization**

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 62.8 kDa. The protein migrates as 70-90 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.**Endotoxin**

Less than 1.0 EU per µg by the LAL method.

**Purity**

&gt;90% as determined by SDS-PAGE.

**Formulation**

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH 7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.***Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

**SDS-PAGE**Influenza B virus [Austria/1359417/2021 (B/Victoria lineage)] HA Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).**Bioactivity-ELISA**

Immobilized Influenza B virus [Austria/1359417/2021 (B/Victoria lineage)] HA Protein, His Tag (Cat. No. HAE-V52H8) at 1 µg/mL (100 µL/well) can bind various dilution ratio of Influenza B Hemagglutinin / HA Antibody, Rabbit IgG (QC tested).

**Discounts, Gifts,  
and more!**



## Background

Neuraminidase (NA) and hemagglutinin (HA) are major membrane glycoproteins found on the surface of influenza virus. Hemagglutinin binds to the sialic acid-containing receptors on the surface of host cells during initial infection and at the end of an infectious cycle. Hemagglutinin also plays a major role in the determination of host range restriction and virulence. As a class I viral fusion protein, hemagglutinin is responsible for penetration of the virus into the cell cytoplasm by mediating the fusion of the membrane of the endocytosed virus particle with the endosomal membrane.

## Clinical and Translational Updates

Discounts, Gifts,  
and more!

